

ART. II.—*Monographia Chalciditum.* By FRANCIS
WALKER, Esq. F.L.S.

(Continued from Vol. I. page 466.)

“ —— the green myriads in the peopled grass.”

Family.—LEUCOPSIDÆ.

GENUS LEUCOPSIS,^a Fabricius.

Vespa . . . Swammerdam, Sulzer, Christ.

Cynips . . . De la Tourette, Gmelin.

*Leucospis. Fabricius, Gmelin, Latreille, Olivier, Villers,
Rossi, Cuvier, Panzer, Jurine, Walckenaer,
Illiger, Spinola, Fuessly, Dalman, Klug,
Fonscolombe.*

Leucopsis. Lamarck, Dumeril.

Corpus punctatum, supra pubescens : caput mediocre, transversum, thorace vix angustius, anticè ubi scapi insident excavatum : oculi mediocres : ocelli 3, supra verticem trigonè dispositi : antennæ maris et fem. similes, medio frontis insertæ, 14-articulatæ, clavatæ, pubescentes ; scapus in canaliculo frontalí receptus ; flagellum subincurvum ; articulus 1^{us}. s. scapus elongatus ; 2^{us}. brevis, cyathiformis ; 3^{us}. elongato-cyathiformis ; 4^{us}. et 7 sequentes ferè lineares, latiores, breviores ; 12^{us}. mediocris, apice angustior ; 13^{us}. et 14^{us}. minimi, vix conspicui : labrum breve, transversum, anticè impressum : mandibulæ ferè rectæ, æquales, bidentatae ; dentes obtusi ; internus brevis, latus, ferè geminus : maxillæ elongatae, graciles, apice acuminatæ, externè pilosæ, internè apicem versus in lobum productæ bipartitum tenuem mollem ciliatum : palpi 4-articulati, filiformes ; articulus 1^{us}. elongato-cyathiformis ; 2^{us}. longior ; 3^{us}. 2^o. brevior ; 4^{us}. 2^o. longior : mentum elongatum, angustum, apice trispinosum : palpi 3-articulati, ferè filiformes, menti apicem versus inserti ; articuli subclavati, 2^{us}. 1^o. brevior, 3^{us}. longior : labium ferè cordiforme, molle ; anguli antici producti ; margo anticus ciliatus, medio impressus : thorax ovatus : prothoracis scutellum maximum, subquadratum, anticè paullò angustius ; segmenta reliqua dorsalia supra occulta ; pectus parvum : mesothoracis scutum maximum ;

^a Λευκὸς albus, ὄψις facies. I think, with Dumeril, that the name of the genus is derived from these words.

parapsides scuto in unum confusæ; scutellum magnum, semi-circulus; paraptera et epimera trigona; sternum magnum, medio canaliculatum: metathoracis scutellum parvum, plerumque apice bispinosum; postscutellum maximum; epimera et paraptera trigona; sternum magnum: propedes^b graciles; coxae trigonæ, mediocres; trochanteres parvi; tibiæ paullò arcuatæ, apice latiores, et dentibus nonnullis minutis et spina elongata valida subarcuata armatae: mesopedes longiores, graciliores; tibiæ rectæ, femoribus paullò longiores, externè basim versus impressæ; spina apicalis brevior, tenuior; cæteri propedem: metapedium coxae maximæ, angulus internus serratus; femora maxima, ovata, externè convexa, internè plana, subtus dentibus plurimis armata; tibiæ valdè arcuatæ, subtus canaliculatae, femoribus appressæ, apicis angulus internus productus, acuminatus; cæteri propedem: tarsi omnes subtus ciliati, articulus 1^{us}. elongatus, sequentes longitudine decrescentes, ultimus 2^o. longior; unguis arcuati, subtus basi dentibus plurimus minutis armati; pulvilli minimi: proalæ plicate; nervus subcostalis costam alæ medium versus attingens, et inde ferè ad apicem productus, ubi costam attingit ramulum emittens furcatum; furca antica mox abrupta; postica alæ apicem versus producta; nervus in alæ disco, brevis, insulatus; nervus 3^{us}. longitudinalis alæ basi proveniens, medio crassior, apicem versus furcatus; furcae ambæ marginem posticum attingentes; nervus 4^{us}. marginis postici medium occupans; areolæ paucæ, apertæ, non benè determinatae: metalæ parvæ; nervus subcostalis a basi, ubi costa jungitur, ferè ad alæ apicem productus; nervi duo longitudinales indistincti; hamus nervo subcostali ante alæ apicem insertus: abdomen sessile, elongato-ovatum, compressum, apice rotundum; *maris*, segmentum 1^{um}. magnum; 2^{um}. maximum, ferè omne abdominis dorsum occupans; 3^{um}. minimum, vix conspicuum; subtus abdomen segmenta 5 ventralia conspicua, subæqualia: *fem.* abdomen dorso ubi oviductus insidet canaliculatum; segmentum 1^{um}. maximum; 2^{um}. minimum, 1^o. ferè occultum; 3^{um}. mediocre; 4^{um}. majus, latera latiora; 5^{um}. parvum; 6^{um}. maximum, 5ⁱ. dorso recurvum, subtus quoque recurvum; subtus abdomen lamina elongata, angusta, carinata, segmenta omnia ventralia occultat: oviductus segmenti 6ⁱ. apice emersus, dorso recurvus.

These insects, in the larva state, are parasitic upon the

^b I have adopted this and similar terms, from Mr. Newman's letter on Osteology, published in Vol. I. p. 400.

mason-bees and wasps. The trophi, particularly the lingua and maxillæ, are more developed than in most *Chalcidites*.

DIVISIO I. *Metafemora subtus dentibus 8 armata.*

Sp. 1. Leuc. grandis.

Leucospis gigas . . . Rossi *Faun. Etrusc.* ed. Illiger. II.
130; Fonsc. *Ann. Sci. Nat.* 26. 273.

Leucospis gallica . . . Rossi *Mant.* 135. 298.

Leucospis dorsigera. Var. Hochenwarth *Schrift. Berl. Ges.*
VI. 341. Taf. 8. fig. 1, 2; Christ.
Bienen. 225. Tab. 19. fig. 9.

Leucospis grandis . . . Klug. *Act. Nat. Cur. Berl.* VI. 66. 1.

Sp. 2. Leuc. gigas. Mas et fem. *Nigra, flava variegata,*
prothoracis scutello flavo marginato, mesoscuti dorso
bimaculato, oviductu abdominis basim non attingente.

Cynips nigra, &c. . . De la Tourette *Acad. des Sci.* 9. 730.
fig. 1—5.

Cynips lugdunæa . . . Gmel. 1. 5. 2653. 26.

Sphex dorsigera . . . Sulzer. *Gesch.* 196. Tab. 27. fig. 11.

Leucospis dorsigera. Fabr. *Syst. Ent.* 361. 1; *Spec. Ins.*
1. 457. 1; *Mant. Ins.* 1. 284. 1;
Gmel. 2739. 1; Oliv. *Encyclop. Hist. Nat. Ins.* VII. 352. 1. Pl. 100.
fig. 8. 9; Fuessly *Archiv.* 2. Tab.
18. 51. fig. 1—10.

Leucospis gallica . . . De Villers *Ent.* III. 261. Tab. 8.
fig. 18.

Leucospis gigas . . . Fabr. *Ent. Syst.* II. 245. 1; *Syst.*
Piezat. 168. 1.; Panz. *Faun. Ins.*
Germ. 84. Tab. 17. 18.; Coqueb.
Illustr. Icon. I. 23. Tab. 6. fig. 1.;
Spin. Ins. Lig. fascic. 1. 63.; Latr.
Hist. Nat. des Ins. XIII. 219. 2.;
Gen. Crust. et Insect. IV. 24.; *Règne*
Anim. III. 475.; *Nouv. Edit.* V.
296.; *Nouv. Dict. d'Hist. Nat.* XVII.
514.; Klug. *Act. Nat. Cur. Berl.*
VI. 66. 2.

*Leucopsis gigas . . . Lam. Anim. sans Vertèbres. IV. 151. ;
Dumeril Dict. des Sci. Nat. XXVI.
169.*

Mas.—Nigra, flavo pubescens: caput inter oculos flavo bimaculatum: labium rufescens: oculi fusci: ocelli sordidè albidi: antennæ apice rufæ; articulus 1^{us}. flavus, extùs et apice supra niger: prothoracis scutellum flavo marginatum; mesothoracis scuti dorsum flavo bimaculatum, latera flavo vittata; super scutelli dorsum macula flava, anticè valdè emarginata; epimera ferè tota flava: metathoracis scutellum apice acutè bispinosum: abdomen flavo bifasciatum et apice maculatum: pedes flavi; pro- et mesocoxæ nigræ; metacoxæ nigræ, flavo supra basi et subtus apice maculatae; trochanteres fusci; pro- et mesofemora basi nigrofusca aut nigra; metafemora nigra, extùs flava nigro maculata; dentes nigri; 1^{us}. brevis; 2^{us}. et 3^{us}. elongati, acuminati; cæteri obtusi, longitudine decrescentes; metatibiæ subtus nigrofuscæ; tarsi fulvi: alæ iridescentes, fuscæ; costa saturatior.

Fem.—Abdomen supra flavo 4-fasciatum, subtus rufofuscum, nitidum; fasciæ dissimiles, interruptæ: abdominis segmenti 1^{mi}. oviductus medium attingens, fuscus; tegmina nigra. (Corp. long. 4½—6 lin.; alar. 9—12 lin.)

July; South of France. M. F. de Laporte has taken it near Paris.

Sp. 3. *Leuc. nigricornis.*^c Fem. *Nigra, flavo-variegata, prothoracis scutello flavo-bifasciato, mesoscuti dorso im-maculato, oviductu abdominis basim non attingente.*

Leucospis dispar. a. Fonscol. Ann. Sci. Nat. 26. 275.

Nigra, flavo pubescens: caput immaculatum: labium rufescens: oculi fusci: ocelli sordidè albidi: antennæ apice fuscae: prothoracis scutellum flavo bifasciatum; fascia antica medio angustior: mesothoracis scuti latera flavo vittata; epimera flavo maculata; metathoracis scutellum inerme?; super dorsum macula flava, anticè valdè emarginata: abdomen utrinque supra flavo 4-maculatum, subtus rufo-fuscum; maculæ dissimiles: abdominis segmenti 1^{mi}. oviductus apicem attingens, fuscus; tegmina nigra: pedes flavi; coxae nigræ; metacoxæ supra basi et subtus apice flavo maculatae; trochanteres nigrofusci; pro- et mesofemora basi nigra; metafemora nigra, externè flava nigro maculata; dentes nigri, 1^{us}. brevis, 2^{us}. et 3^{us}. elongati, acumi-

^c This, and many of the following species, were sent to me by M. F. de Laporte, with the MS. names which I have adopted.

nati; cæteri obtusi, longitudine decrescentes; metatibiae subtus nigro-fuscæ; tarsi fulvi: alæ iridescentes, fuscæ; costa saturatior. (Corp. long. 6 lin.; alar. 12 lin.)

Taken near Paris, by M. F. de Laporte.

Sp. 4. Leuc. intermedia. Fem. *Nigra, flavo variegata, prothoracis scutello flavo-marginato, mesoscuti dorso unimaculato, oviductu abdominis basim superante.*

Leucospis dorsigera . *Rossi Mant.* 134. 297. *De Villers Ent.* III. Pl. 8, fig. 17. *Panz. Faun. Ins. Germ.* 15. Tab. 17.
Leucospis intermedia. *Illiger. Edit. Faun. Etrusc.* II. 130. *Klug Act. Nat. Cur. Berl.* 67. 4.

Nigra, flavo pubescens: caput inter oculos flavo bimaculatum: labium rufescens: oculi fuscæ: ocelli sordidè albidi: antennæ nigrae; articulus 1^{us}. flavus, extùs et apice supra niger: prothoracis scutellum flavo marginatum: mesothoracis scuti dorsum flavo unimaculatum, latera flavo vittata; super scutelli dorsum macula flava, anticè valdè emarginata; epimera flavo maculata: metathoracis scutellum apice acutè bispinosum: abdomen flavo 4-fasciatum, subtus rufo-fuscum; fasciæ dissimiles, basalis et apicalis interruptæ: oviductus mesothoracis ad scutelli medium productus, fuscus; tegmina nigra: pedes flavi; coxae nigrae; metacoxæ supra basi flavo maculatae; trochanteres fuscæ; pro- et mesofemora basi nigra; metafemora nigra, extùs flava, nigro maculata; metatibiæ subtus nigro-fuscæ; tarsi fulvi: alæ iridescentes, fuscæ; costa saturatior. (Corp. long. 5 lin.; alar. 9 lin.)

July; South of France.

Sp. 5. Leuc. varia. Fem. *Nigra, flavo variegata, prothoracis scutello flavo bifasciato, mesoscuti dorso bimaculato, oviductu abdominis basim non attigente.*

Leucopsis intermedia . *Lamarck. Anim. sans Vertèbres.* IV. 152.

Leucospis varia . . *Klug Act. Nat. Cur. Berl.* VI. 67. 3.

Nigra, flavo pubescens: caput inter oculos flavo bimaculatum: labium rufescens: oculi fuscæ: ocelli sordidè albidi: antennæ NO. I. VOL. II. D

apice rufo-fuscæ ; articulus 1^{us}. fulvus, apice fuscus ; 2^{us}. nigro-fuscus ; 3^{us}. et 4^{us}. rufi : prothoracis scutellum flavo bifasciatum ; fascia antica medio angustior, posticè incisa : mesothoracis scuti latera flavo vittata, dorsum flavo bimaculatum ; super scutelli dorsum macula flava, anticè valde emarginata ; epimera ferè tota flava : metathoracis scutellum apice acutè bispinosum : abdomen supra flavo 4-fasciatum, subtus fuscum ; fasciæ dissimiles, interruptæ : abdominis segmenti 1^{mi}. oviductus ferè basim attingens, fuscus ; tegmina nigra : pedes fulvi ; coxæ nigræ ; metacoxæ supra basi et subtus apice flavo maculatæ ; trochanteres nigri ; pro- et mesofemora basi fusca ; metafemora utrinque nigro maculata ; dentes nigri ; 1^{us}. brevis ; 2^{us}. et 3^{us}. elongati, acuminati ; sequentes obtusi, longitudine decrescentes : alæ iridescentes, fuscæ ; costa saturatior. (Corp. long. 4—4½ lin. ; alar. 9—9½ lin.)

July ; South of France.

Sp. 6. Leuc. aculeata.

Leucospis aculeata . *Klug. Act. Nat. Cur. Berl.* VI. 68. 5.

DIVISIO II. *Metafemora subtus dentibus 10 armata.*

Sp. 7. Leuc. Biguetina. Mas et Fem. *Nigra, flavo variegata, prothoracis scutello flavo bifasciato, fascia postica abbreviata.*

Leucospis gibba . . *Klug. Act. Nat. Cur. Berl.* VI. 70. 8.

Leucospis Biguetina *Jurine, Nouv. Method. Hyménopt. &c.* 307. Pl. 13. 45.

Mas. Nigra, pubescens : caput immaculatum : labium rufescens : oculi fusi : ocelli sordidè albidi : antennæ omnino nigræ : prothoracis scutellum flavo bifasciatum ; fascia antica medio angustior, postica abbreviata : mesothoracis scuti latera flavo vittata ; super scutelli dorsum fascia flava, anticè valde emarginata ; epimera ferè tota flava : metathoracis scutellum apice acutè bispinosum : abdomen gibbum, supra flavo trifasciatum : pedes flavi ; coxæ nigræ ; trochanteres fusi ; pro- et mesofemora basi fusca ; metafemora nigra, apice supra flava, subtus dentibus 10 nigris armata, dens basalis magnus, cæteri minimi ; metatibiæ subtus pallidè fuscæ ; tarsi fulvi : alæ iridescentes, subfuscæ ; costa et apex saturiores.

Fem. Abdomen flavo 4-fasciatum; fasciæ 3^a. et 4^a. interruptæ, hæc quoque abbreviata: oviductus abdominis medium attingens.
(Corp. long. 3½ lin.; alar. 6½ lin.)

Var. β.—*Fem.* Mesothoracis scuti dorsum flavo bimaculatum.

Described from specimens taken near Paris by M. F. de Laporte.

DIVISIO III. *Metafemora subtus dentibus 15 armata.*

Sp. 8. Leuc. dorsigera. Mas et Fem. *Nigra, flavo variegata, prothoracis scutello flavo bifasciato, fascia antica abbreviata.*

Leucospis cœlogaster . Schrank *Schriften Berl. Gesellschaft.* 1. 301. Tab. 8. fig. 4—6. *Faun Boica* II. 2. 221. 1980. *Gmel.* 2741. 2.

Oliv. Encycl. Hist. Nat. Ins. Tom. VII. 532. 3. *Hochenwarth. Schriften Berl. Ges.* VI. 344. Tab. 8. fig. 34.

Leucospis dubia . . Schrank. *Fauna Boica.* II. 2. 222. 1981.

Leucospis dorsigera . Fabr. *Ent. Syst.* 2. 246. 2.; *Syst. Piezat.* 168. 2. *Latr. Hist. Nat. des Ins.* XIII. 281. 1. *Gen. Crust. et Insect.* IV. 24. *Règne Anim.* III. 475; *Nouv. Edit.* V. 296. *Nouv. Dict. d'Hist. Nat.* XVII. 513. Pl. G. 3. 7. *Panz. Faun. Ins. Germ.* 58. 15. *Illig. Edit. Faun. Etrusc.* II. 130. et 280. 856. β. *Spin. Ins. Lig.* Fasc. I. 63. *De Vill. Ent.* III. 260. 1. Tab. 8, fig. 17. *Klug Act. Nat. Cur. Berl.* 69. 6.

Leucospis dispar . . Fabr. *Syst. Piezat.* 169. 6.

Leucospis intermedia *Spin. Ins. Lig.* Fascic 4. 236. *Fonscol. Ann. Sci. Nat.* 26. 274.

Leucopsis dorsigera . Lam. *Anim. sans Vertèbres.* IV. 151. Dumeril. *Dict. des Sci. Nat.* 26. 169. Pl. 34, fig. 2.

Mas. Nigra, pubescens: labium rufescens: oculi fusci: ocelli sordidè albidi: antennæ nigræ; articulus 1^{us}. flavus, apice et supra fuscus:

prothoracis scutellum flavo bifasciatum; fascia antica abbreviata; postica per scutelli latera producta: mesothoracis scutum immaculatum; super scutelli dorsum fascia flava integra aut paullò emarginata; epimera flavo maculata: metathoracis scutellum inerme: abdomen supra flavo 4-fasciatum; fasciae 1^{ma}. et 4^{ta}. abbreviatæ: pedes flavi; coxae nigræ; metacoxæ subtus apice flavo maculatæ; trochanteres nigro-fusci; pro- et mesofemora basi fusca; metafemora nigra, subtus versus basin et supra flava; dentes nigri, basalis magnus, cæteri minimi; metatibiæ subtus nigræ; tarsi fulvi: alæ iridescentes, subfuscæ; costa et apex saturiores.

Fem. Scapi omnino flavi: abdomen supra flavo trifasciatum, subtus rufo-fuscum; fasciae interruptæ, inter 1^{am}. et 2^{am}. maculæ duæ laterales flavæ: oviductus abdominis basim attingens: profemora nigro-fusca, apice flava; intermedia pallidiora: protibiæ extùs fusco vittatæ. (Corp. long. 2 $\frac{2}{3}$ —4 lin.; alar. 4—6 lin.)

Var. β.—Mas. Antennæ omnino nigræ: prothoracis fascia postica non per scutelli latera producta: metacoxæ nigræ, apice subtus fusco maculatæ; pro- et mesofemora nigro-fusca, apice flava; metafemora nigra, basi subtus et supra apice flavo maculata; pro- et mesotibiæ extùs fusco vittatæ; metatibiæ subtus et intùs nigræ.—Species distincta?

Var. γ.—Fem. Abdominis fascia apicalis angusta, abbreviata: metafemora nigra, supra apice flava.

Described from specimens taken near Paris, by M. F. de Laporte.

Sp. 9. Leuc. bifasciata.

Leucospis, &c. . . *Fuessly Archiv.* III. Tab. 18. fig. 11.
Leucospis bifasciata . *Klug. Act. Nat. Cur. Berl.* 70. 7.

Family CHALCIDIDÆ.

Structura varia: reliquis ordinis characteres sequentes immutati eam distinguunt; oviductus infra abdomen occultus; metapedes femoribus et coxis maximis, tibiis arcuatis; proalæ ultra medium nervus ordinarius ramulum emittens brevissimum, nonnunquam furcatum: ocelli 3, supra verticem trigonè dispositi: pro- et mesocoxæ trigonæ, mediocres: tarsi plerumque graciles; articulus 1^{us}. elongatus; sequentes longitudine decrescentes; ultimus 2^o longior.

Characteres Generum.

Caput	inerme. Petiolus	elonga- tus. Antennæ	{ medio frontis insertæ prope os insertæ	I. SMIERA.
		brevissi- mus. Antennæ		II. EPITRANUS.
	armatum	medio frontis insertæ	{ acuminatæ non acu- minatae. Metatibiae apice	III. CHALCIS.
		prope os insertæ. Metatibiae apice		IV. HALTICELLA.
		graciles Metatarsi crassi		V. HOCKERIA. VI. NOTASPIS.
				VII. DIRHINUS.

GENUS I.—SMIERA, *Spinola.*

Sphex. *Linnæus, Fabricius, Villers, Sulzer, Schrank, Christ.*

Chrysis. *Fabricius.*

Vespa. *Geoffroy, Fourcroy.*

Chalcis. *Fabricius, Gmelin, Panzer, Hubner, Rossi, Latreille, Stewart, Olivier, Spinola, Donovan, Lamarck, Leach, Samouelle, Dalman, Fonscolombe, Cuvier, Jurine, Illiger, Walckenaer, Dumeril.*

Smiera. *Spinola, Curtis.*

Corpus punctatum, pubescens : caput mediocre, transversum, thorace vix angustius, anticè ubi scapi insident excavatum : antennæ 13-articulatae, medio frontis insertæ, *maris* fusiformes aut filiformes, *fem.* clavatae ; articulus 1^{us}. elongatus, in-canaliculo frontali receptus ; 2^{us}. minutus ; 3^{us}. minimus, vix conspicuus ; 4^{us}. elongatus ; 5^{us}. et 5 sequentes longitudine decrescentes ; clava triarticulata, ovata : labrum breve, transversum, apice ciliatum : mandibulæ parvæ ; una ferè recta, subquadrata, tridentata ; altera arcuata, bidentata ; dentes obtusi : maxillæ elongatae, apicem versus internè in lobum productæ latum ciliatum apice incisum ; palpi 4-articulati, filiformes ; articuli 1^{us}. et 3^{us}. breviores ; 2^{us}. et 4^{us}. longiores : mentum elongatum, angustum ; palpi 3-articulati, subfiliformes, articulus 3^{us}. acuminatus, setosus : labium angustum, fissum : thorax ovatus : prothoracis scutellum minimum, posticè incurvum ; pectus parvum : mesothoracis scutum maximum ; parapsides optimè determinatae ; scutellum maximum, semicirculus ; paraptera et epimera trigona, maxima ; sternum parvum : metathoracis scutellum parvum ;

postscutellum maximum; paraptera et epimera trigona; sternum parvum: petiolus elongatus, linearis, abdomine brevior: abdomen glabrum, gibbum, compressum, coarctatum, nonnunquam apice elongatum et acuminatum; segmentum 1^{um.}^d maximum; cætera breviora; segmenta ventralia pauca *maris* abdominalis apicem versus conspicua: fem. lamina angusta segmenta omnia ventralia et oviductum nisi ad apicem abscondit: propedes breves; trochanteres parvi; femora subclavata; tibiae apice spina elongata valida armata; unguis elongati, subarcuati: pulvilli minutus: mesopedes longiores et graciliores, cætera propedem: metapедum coxae maximæ, elongatae; femora maxima, ovata, subtus dentibus plurimus armata, quorum basalis maxima; tibiae validè arcuatæ, subtus canaliculatae, femoribus appressæ; apicis angulus internus productus, acuminatus; cætera propedem: proalæ nervus ordinarius ante medium costæ junctus, inde per costam ferè ad alæ apicem productus: nervi duo longitudinales sinuati, indistincti, anticus furcam emittens ramulo stigmaticalij junctam: metalæ angustæ; nervus ordinarius ultra medium productus, apice hamo armatus.

DIVISIO 1. *Abdomen breve, vix longius quam latum.*

Latreille supposed that these insects infest the *Stratiomydæ*, or other *Diptera*, that are aquatic in their larva state. *C. xanthostigma*, Dalm. is parasitic upon a species of *Hylotoma*.

Sp. 1. Smi. nigrifex. Mas et Fem. *Nigra, petiolo et pedibus flavis, his nigro variegatis.*

Sphex nigrifex. . . Sulzer, *Hist. Ins.* 191. I. fig. 27. 1.

Vespa &c. . . . Geoffroy, *Ins. Par.* II. 380. 16.

Vespa dearticulata. Fourc. *Ent. Par.* Tom. II. 437. 16.

Sphex sispes . . . Fabr. *Sp. Ins.* I. 446. 61. Villers, *Ent.* Tom. I. 222. 6.

Chrysis sispes . . . Fabr. *Syst. Ent.* 359. 15.

Chalcis sispes . . . Fabr. *Mant. Ins.* I. 272. 1. *Ent. Syst.*

II. 194. 1. *Syst. Piezat.* 159. 1.

Gmel. *Syst. Nat.* I. 5. 2742. 1.

Hubn. *Naturf.* 24. 54. 18. Tab. 2.

^d The second segment of my preceding descriptions; the petiole being considered as the first. The other alterations require no explanation.

fig. 22. *Panz.* 77. 11. *Stew.* II.
 236. *Rossi, Faun. Etrusc.* II. 58.
 802. *Oliv. Encycl. Méthod.* V. 438.
 2. *Spin. Ins. Lig.* fasc. 1. 62.
Latr. Règne Anim. III. 474; *Nouv. Edit.* V. 295. *Nouv. Dict. d'Hist. Nat.* VI. 23. 10. *Lam. Anim. sans Vertèbres,* IV. 153. *Fonscol. Ann. Sci. Nat.* XXVI. 275. 1.

Smiera sispes . . . Spinola. Ann. du Muséum d'Hist. Nat. Tom. XVII.

Smiera petiolatus . Curtis Brit. Ent. 472.

Mas.—*Nigra*, pilis canis vestita : caput inter oculos flavo bimaculatum : oculi ocellique fusci : antennæ fusiformes, corporis dimidio longiores ; scapus nitidus ; clava articulis 2 præcedentibus paullò brevior : palpi maxillares articulis 4 ; 1^{us}. medioeris ; 2^{us}. vix longior ; 3^{us}. brevior ; 4^{us}. elongatus, fusiformis, setosus : palpi labiales articulis 3 ; 1^{us}. et 2^{us}. breves ; 3^{us}. longior : mesothoracis sentellum apice bisponosum ; squamulae flavæ : petiolus flavus : abdomen nitidum ; segmenta 1^{um}. et 2^{um}. maxima ; sequentia parva, subæqualia : pedes nigri ; metacoxæ supra apice spinosæ ; pro et mesofemora apice flava ; metafemora flava, apice et supra basi nigra, subtus dentibus 13 nigris armata ; protet mesotibiæ fuscæ, apice basique flavæ ; metatibiæ apice flavæ ; tarsi fulvi, apice fusci : alæ subfuscæ ; nervi fusci.

Fem.—Antennæ breviores, tenuiores, clavatae : metafemorum dens basalis multò longior et crassior : petiolus brevior : abdominis segmenti 1ⁱ. latera latiora ; 2^{um}. parvum, 1ⁱ. margine ferè occultum ; 3^{um}. magnum, latera 1ⁱ. margine ferè occulta ; sequentia minima : oviductus rufo-fuscus. (Corp. long. 3 lin. ; alar. 5 lin.)

Common in the South of Europe ; rare in the North. July ; South of France ; on aquatic plants. M. F. de Laporte has taken it near Paris.

Sp. 2. Smi. sispes. Mas et. Fem. *Nigra, metafemoribus tarsisque rufis.*

Sphex sispes . . *Linn. Syst. Nat.* XII. 2. 943. 13. *Faun. Suec.* 1657.

Chalcis clavipes. *Fabr. Maut. Ins.* 1. 272. 2. *Ent. Syst.* III. 2. 195. 2. *Syst. Piezat.* 159. 2. *Gmel. Syst. Nat.* I. 5. 2742. 2. *Hubn.*

Naturf. 24. 56. 19. *Tab.* 2. *fig.* 23.
Oliv. *Encycl. Méthod.* V. 438. *Rossi*
Faun. Etrus. II. 58. 803. *Latr. Règne*
Anim. III. 474; *Nouv. Edit.* V. 295.
Nouv. Dict. d'Hist. Nat. VI. 13. *Pl. B.*
23. 10. *Panz.* 78. *fig.* 15. *Don.* XI.
57. 379. *Lam. Anim. sans Vertèbres,*
IV. 153. *Leach, Ed. Encycl.* IX. 144.
Samouelle, 271. *Pl. 8. fig. 6.* *Fonscol.*
Ann. Sci. Nat. XXVI. 276. 2.

Smiera clavipes . *Spin. Ann. du Muséum, &c.* Tom. XVII.
Chalcis sispes . *Dalm. Act. Kongl. Vetensk. Acad. Handl.*
für är. 1820.

Smiera sispes . . *Curtis, Brit. Ent.* 472.

Mas.—*Nigra*, pilis canis vestita: oculi ocellique fuscæ: antennæ subfiliformes, corporis dimidio breviores; scapus nitidus: palpi maxillares articulis 4; 1^{us}. brevis, extùs excavatus; 2^{us}. multò longior; 3^{us}. brevis, 1^o. æqualis: mentum quàm *S. nigrifex* latius: palpi articulis 3 ferè æqualibus; 2^{us}. paullò brevior: mesothoracis scutellum apice bisponosum; squamulæ fuscæ: petiolus quàm *S. nigrifex* brevior: abdomen nitidum: pedes nigri; pro- et mesofemora apice rufa; metafemora rufa, apice nigra, subtus dentibus 8 inæqualibus armata; pro- et mesotibiæ nigro-fuscæ; metatibiæ nigrae; tarsi rufi, apice fuscæ; alæ subfuscæ, apice fuscæ; nervi fuscæ.

Fem.—Antennæ paullò breviores, clavatæ: oviductus rufo-fuscus.
(Corp. long. 2 $\frac{3}{4}$ —3 $\frac{1}{4}$ lin.; alar. 4 $\frac{3}{4}$ —5 $\frac{1}{4}$ lin.)

Var. β.—Metafemora extùs flavo maculata.

Described from specimens taken at Paris by M. F. de Laporte. Unlike *C. nigrifex*, it abounds more in the North than in the South of Europe.

There are three more described European species; viz.—

Sp. 3. Smi. melanaris.

Chalcis melanaris . . *Dalm. Act. Kongl. Vetensk. Acad. Handl. für är.* 1820.

Smiera Macleanii . . *Curtis, Brit. Ent.* 472.

Taken lately in Essex.

Sp. 4. Smi. biguttata.

Chalcis biguttata . . . *Spin. Ins. Lig.* Fasc. 4^{us}. 231.

Sp. 5. Smi. xanthostigma.

Chalcis xanthostigma. *Dalm. Act. Kongl. Vetensk. Acad. Handl. für är.* 1820.

DIVISIO II. *Abdomen apice elongatum et acuminatum.*

The exotic species of this division, and of some genera nearly allied to it, are very numerous.

Sp. 6. Smi. subpunctata. Mas et Fem. *Flava, nigro variegata, alis hyalinis.*

Mas.—Flava, vix pubescens : caput subtus nigro fasciatum : oculi virides : ocelli fuscii : antennæ subfusciformes, fuscæ, subtus fulvæ : mesothoracis scutum anticè nigrum, medio nigro vittatum ; parapides nigro maculatae ; paraptera nigro maculata, maculæ inter scutum et scutellum conjunctæ ; super scutellum macula teli-formis nigra ; pectus nigro bimaculatum : metathoracis scutellum nigro fasciatum : petiolus abdominis dimidio brevior : abdomen elongato-ovatum, nitidum, fulvum ; maculæ 5 dorsales et apex nigræ ; segmentum 1^{um}. flavum, magnum ; cætera parva ; pedes flavi ; metacoxæ extùs nigro maculatae ; metafemora subtus basi et apice nigro maculata, subtus dentibus 14 nigris armata ; dens basalis magna, cæteri minimi ; metatibiæ basi, apice et subtus fuscæ ; unguis et pulvilli fuscii : alæ hyalinæ, iridescentes ; nervi fulvi.

Fem.—Antennæ subclavatae : abdomen apice acuminatum ; segmentum 1^{um}. magnum ; 2^{um}. et 4 sequentia minuta ; cætera elongata : oviductus fuscus. (Corp. long. 2½ lin.; alar. 3 lin.)

Var. β—*Mas*, abdomen fulvum, fusco fasciatum.

Taken in St. Vincent's island, by the Rev. Lansdown Guilding : described also from a Cayenne specimen in the collection of M. F. de Laporte.

Sp. 7. Smi. fulvescens. Mas et Fem. *Lætè ferruginea, abdomine fusco, alis hyalinis.*

Mas.—Ferruginea, subtus pallidior, vix pubescens: oculi virides: ocelli fuscæ: antennæ subfusciformes, fuscæ, subtus fulvæ: petiolus gracilis, abdominis dimidio brevior: abdomen elongato-ovatum, nitidum, ferrugineum, supra fusco fasciatum: pedes flavescentes; metafemora subtus dentibus 14 nigris armata; dens basalis magna, cæteri minimi; metatibiæ subtus fuscæ; unguis et pulvilli fuscæ: alæ hyalinæ, iridescentes; nervi fulvi.

Fem.—Antennæ subclavatae: petiolus quam maris brevior: abdomen acuminatum; segmenta apicalia elongata; apex nigro-fusca: oviductus omnino occultus. (Corp. long. $2\frac{1}{2}$ lin.; alar. 3 lin.)

Var. β.—*Mas*, abdomen fuscum, basi fulvo fasciatum.

Taken in St. Vincent's island, by the Rev. Lansdown Guilding.

GENUS II.—EPITRANUS,^e *Walker.*

Fem.—Corpus punctatum, sparsè pubescens: caput mediocre, transversum, ferè trigonum: antennæ 14-articulatae, subfusciformes, corporis dimidio breviores, basi approximatæ, prope os insertæ: articulus 1^{us}. valde elongatus, flagelli dimidio longior; 2^{us}. brevis; 3^{us}. minimus; 4^{us}. et 11 sequentes subæquales; clava 3-articulata, conica, articulis 10°. et 11°. brevior et angustior: mandibulæ angustæ; una recta, bidentata, dentes acuti, externus magnus; altera apice arcuata, dente brevi acuto terminata: thorax ovatus: prothoracis scutellum mediocre, posticè incurvum; pectus parvum: mesothoracis scutum mediocre; parapsides optimè determinatæ, maximæ, convexæ; paraptera et epimera trigona, magna; scutellum maximum, ferè rotundum; sternum parvum: metathoracis scutellum et postscutellum parva; sternum magnum: petiolus linearis, abdominis dimidio longior: abdomen elongato-ovatum, vix compressum, subtus carinatum, apice acuminatum; segmentum 1^{um}. maximum, ferè ad apicem productum; cætera minima, brevissima: oviductus occultus: pro- et mesopedum coxæ parvæ; femora clavata; tibiæ apice spina elongata, valida, arcuata armata; unguis et pulvilli minimi: metapedum coxæ apice angustiores; femora subtus dentibus 9 armata; quorum basalis maximus, obtusus; cæteri minimi, acuti; tibiæ *Smieræ*: alæ breves; nervi indistincti.

Sp. 1. Epi. fulvescens. Fem. *Rufo-fuscus, tarsis flavis, alis albis.*

^e ἐπὶ antē, τραυός planus.

Caput punctatum : oculi ocellique fusi : antennæ fuscae ; articulus 1^{us}. fulvus : thorax punctatus ; squamulae flavæ : petiolus striatus : abdomen nitidum, glabrum : metatibiæ subtus fuscæ ; tarsi flavi ; ungues et pulvilli fusi : alæ albæ, iridescentes ; nervi flavi. (Corp. long. 1½—2 lin. ; alar. 2—2½ lin.)

Taken in St. Vincent's Island, by the Rev. Lansdown Guilding.

GENUS III.—CHALCIS, *Fabricius*.

Vespa *Limnæus*, *Geoffroy*, *Fourcroy*.

Chalcis . . . *Fabricius*, *Gmelin*, *Rossi*, *Latreille*, *Panzer*,
Olivier, *Lamarck*, *Cuvier*, *Spinola*, *Dumeril*,
Dalman, *Leach*.

Brachymeria. *Westwood*.

Corpus punctatum, pubescens : caput mediocre, breve, transversum, thorace non latius, anticè ubi scapi insident excavatum : oculi mediocres : ocelli trigonè dispositi : antennæ 13-articulatae, plus minusve fusiformes, medio frontis insertæ ; articulus 1^{us}. elongatus ; 2^{us}. mediocris ; 3^{us}. minimus ; 4^{us}. et 6 sequentes crassi, subæquales, longitudine decrescentes ; clava 3-articulata, conica, articulis 2 præcedentibus brevior : mandibulæ arcuatæ, bidentatae ; dentes obtusi, internus brevior : maxillæ elongatae, angustæ, intùs apicem versus in lobum tripartitum productæ : palpi 4-articulati, ferè filiformes ; articuli 1^{us}. et 3^{us}. breves ; 2^{us}. longior ; 4^{us}. multò longior, fusiformis : mentum ovatum : palpi articulis 3 ; 1^{us}. et 3^{us}. apice crassiores ; 2^{us}. brevis : labium angustum, fissum : thorax ovatus : prothoracis scutellum magnum, subquadratum, posticè excavatum, anticè angustius ; pectus parvum : mesothoracis scutum maximum ; parapsides benè determinatae ; scutellum maximum, latius quām longum, apice plerumque bispinosum ; paraptera et epimera magna, trigona ; sternum parvum : metathorax parvus ; scutellum mediocre : petiolus brevissimus : *maris* abdomen gibbum, vix duplo longius quām latum ; segmentum 1^{um}. maximum, dimidium occupans ; 2^{um}. mediocre ; sequentia minima ; segmenta 7 aut plura ventralia subtus abdomen conspicua, quorum 1^{um}. et 2^{um}. magna, reliquæ minuta : *fem.* abdomen paullò longius et acutius, infra carinatum ; segmenta ventralia vix conspicua : propedes mediocres ; coxae trigonæ ; trochanteres parvi ; femora paullò incrassata ; tibiae subclavatae, reætæ, apice spina magna arcuata armatae ; ungues et

pulvilli mediocres: mesopedes paullò longiores et tenuiores; spina tibialis brevior et gracilior; cætera propedum: metapedes et alæ *Smieræ*.

Sp. 1. Chal. femorata. Mas. *Nigra, pedibus flavis, metafemoribus nigro fasciatis.*

Chalcis femorata . *Panz. Faun. Ins. Germ.* Fasc. 84. fig. 16.
Chalcis flavipes . *Latr. Gen. Crust. et Insect.* IV. 26.

Nigra, pilis canis vestita: oculi fusci: ocelli rufo-fusci: antennæ subfusciformes, nigræ, apice rufo-fuscae: mesothoracis scutellum apice acutè bispinosum; squamulæ flavæ: abdomen nitidum, punctatum: pedes flavi; coxae nigræ; trochanteres fusci; pro- et mesofemora basi nigra; metafemora nigro cingulata, subtus dentibus 12 inæqualibus armata; pro- et mesotibiae subtus fusco maculatae; metatibiarum canaliculorum margines fusci; ungues et pulvilli fusci: alæ subhyalinæ; nervi fusci. (Corp. long. $2\frac{1}{2}$ —3 lin.; alar. $4\frac{1}{2}$ —5 lin.)

Taken near Paris, from the pupæ of *Zygæna Filipendulæ*, by M. F. de Laporte.

Sp. 2. Chal. flavipes. Fem. *Nigra, pedibus flavis, metafemoribus basi nigris, alis subhyalinis.*

Chalcis flavipes . *Fabr. Ent. Syst.* II. 197. 10. *Syst. Piezat.*
 167. 32. *Latr. Hist. Nat. des Ins.* XIII.
 220. *Panz. Fasc.* 78, fig. 16. *Fonscol. Ann. Sci. Nat.* XXVI. 276.

Nigra, pilis canis vestita: oculi fusci: ocelli rufo-fusci: antennæ fusiformes, nigræ, apice rufo-fuscae: mesothoracis scutellum inerme; squamulæ flavæ: abdomen nitidum, ferè glabrum: pedes flavi; coxae nigræ; trochanteres fusci; pro- et mesofemora basi nigra; metafemora nigra, apice flava, subtus dentibus 13 aut 14 inæqualibus armata; pro- et mesotibiae subtus fusco maculatae; metatibiae subtus fuscae; ungues et pulvilli fusci: alæ subhyalinæ; nervi fusci. (Corp. long. 3 lin.; alar. 5 lin.)

Taken near Paris, by M. F. de Laporte. It may be the female of the preceding species.

Sp. 3. Chal. distinguenda. Mas et Fem. *Præcedenti,*
minor, nigra, pedibus flavis, metafemoribus basi nigris, alis
hyalinis.

Mas. Nigra, pilis canis vestita : oculi fusci : ocelli rufo-fusci : antennæ subfusciformes, nigræ, apice rufo-fuscæ : mesothoracis scutellum dentibus 2 brevissimis obtusis armatum ; squamulæ flavæ : abdomen nitidum, sparsè punctatum : pedes flavi ; coxæ nigræ ; trochanteres fusci ; femora nigra, apice flava ; metafemora subtus dentibus 12 armata ; pro- et mesotibiale subtus fusco maculatae ; metatibiae subtus fuscæ ; unguis et pulvilli fusci : alæ hyalinæ ; nervi fusci.

Fem. Antennæ fusiformes, paullò breviores et crassiores : mesothoracis scutellum inerme. (Corp. long. 2—2½ lin. ; alar. 4—4½ lin.)

Taken near Paris, by M. F. de Laporte. July; South of France.

Sp. 4. Chal. tibialis. *Mas.* *Nigra, pedibus flavis, pro- et mesotibiis nigro maculatis.*

Nigra, pilis canis vestita : oculi fusci : ocelli rufo-fusci : antennæ nigræ, apice rufo-fuscæ : mesothoracis scutellum apice bispinosum ; squamulæ flavæ : abdomen nitidum, sparsè punctatum et pubescens : pedes flavi ; coxæ nigræ ; trochanteres fusci ; femora nigra, apice flava ; metafemora subtus dentibus 12 armata ; pro- et mesotibiale extùs et subtus nigro maculatae ; metatibiae subtus nigræ ; unguis et pulvilli fusci : alæ hyalinæ ; nervi fusci. (Corp. long. 2 lin. ; alar. 4 lin.)

Taken near Paris, by M. F. de Laporte.

Sp. 5. Chal. annulipes. *Mas et Fem.* *Nigra, pedibus flavis, tibiis omnibus nigro maculatis.*

Chalcis flavipes . . *Fabr. Ent. Syst. II.* 197. 10. *Syst. Piezat.*
 167. 32.?

Mas. Nigra, pilis canis vestita : oculi fusci : ocelli rufo-fusci : antennæ nigræ, apice rufo-fuscæ : mesothoracis scutellum apice tubercu-

latum; squamulæ flavæ: abdomen nitidum, sparsè punctatum et pubescens: pedes flavi; coxæ nigræ; trochanteres fusi; pro- et mesofemora basi nigra; metafemora nigra, apice supra flavo maculata, subtus dentibus 12? armata; pro- et mesotibiæ extùs et subtus nigro maculatæ; metatibiæ basi et medio nigro-fuscæ, subtus quoque nigro-fuscæ; ungues et pulvilli fusi: alæ hyalinæ; nervi fusi.

Fem. Antennæ paullò breviores et crassiores: scutellum inerme. (Corp. long. $1\frac{1}{4}$ — $2\frac{1}{2}$ lin.; alar. $2\frac{1}{4}$ — $4\frac{1}{2}$ lin.)

Described from specimens taken in St. Vincent's Island, by the Rev. Lansdown Guilding: also, from a Cayenne one in the collection of M. de Laporte.

Sp. 6. Chal. cingulata. Fem. *Nigra, pedibus flavis, tibiis omnibus fusco maculatis.*

Nigra, pilis canis vestita: oculi fusi: ocelli rufo-fusi: antennæ nigræ, apice rufo-fuscæ: mesothoracis scutellum inerme; squamulæ flavæ: abdomen nitidum, ferè glabrum, sparsè pubescens: pedes flavi; coxæ nigræ; trochanteres fusi; femora nigra, apice flava; metafemora subtus dentibus 12 armata; pro- et mesotibiæ extùs et subtus fusco maculatæ; metatibiæ supra medio et subtus omnino fuscæ; ungues et pulvilli fusi: alæ hyalinæ; nervi fusi. (Corp. long. 2— $2\frac{1}{2}$ lin.; alar. 4— $4\frac{1}{2}$ lin.)

Taken near Paris, by M. F. de Laporte.

Sp. 7. Chal. minuta. Mas. et Fem. *Nigra, pedibus flavis nigro variegatis, tarsis rufis.*

- Vespa minuta . . . *Linn. Syst. Nat.* 952. 28.
- Vespa, &c. . . . *Geoffroy. Ins.* II. 380. 15.
- Vespa femoralis . . . *Fourc. Ent. Par.* II. 437. 15.
- Chalcis minuta . . . *Fabr. Mant. Ins.* I. 272. 3. *Ent. Syst.* II. 195. 4. *Syst. Piezat.* 165. 23. *Gmel. Syst. Nat.* I. 5. 2742. 3. *Latr. Hist. Nat. des Ins.* XIII. 220. *Règne Anim.* III. 474; *Nouv. Edit.* V. 296. *Nouv. Dict. d'Hist.*

Nat. VI. 13. *Rossi, Faun. Etrusc.*
II. 58. 804. *Oliv. Encycl. Méthod.*
V. 439. 5. *Panz. Fasc. 32. Tab. 6.*
Lam. Anim. sans Verteb. IV. 153.
Dum. Dict. des Sci. Nat. VIII. 69.
Pl. 34. fig. 1. *Leach, Edin. Encycl.*
IX. 144. *Fonscol. Ann. Sci. Nat.*
XXVI. 277.

Chalcis femorata . *Dalm. Kongl. Vetens. Acad. für är*
1820.

Brachymeria minuta. *Westw. Lond. & Edinb. Phil. Mag.*
Third Series. Vol. I. 127.

Mas. Nigra, pilis canis vestita: oculi fusci: ocelli rufo-fusci: antennæ nigræ, apice rufo-fuscae: mesothoracis scutellum apice bispinosum; squamulae flavæ: abdomen ferè glabrum, sparsè pubescens: pedes flavi; coxae nigræ; trochanteres fusci; femora nigra, apice flava; metafemora subtus dentibus 12 armata; tibiae fusco cingulatæ; metatibiæ subtus quoque fuscae, basi rufo-fuscae; tarsi pallidè rufi; unguis et pulvilli fusci: alæ subfuscæ; nervi fusci.

Fem. Antennæ paullò breviores et crassiores: scutellum inerme: abdomen longius. (Corp. long. $2-2\frac{1}{2}$ lin.; alar. $4-4\frac{1}{2}$ lin.)

Var. β.—Mas. Tibiae nigro cingulatæ.

Taken near Paris, by M. F. de Laporte. July; South of France. September; Lizard Point, Cornwall.

Sp. 8. Chal. podagrca. Fem. *Nigra, pedibus rufis, flavo variegatis.*

Chalcis podagrca. *Fabr. Mant. Ins. I.* 272. 5. *Ent. Syst.*
II. 196. 6. *Syst. Piezat.* 166. 24.
Gmel. Syst. Nat. 1. 5. 2743. 5. *Hubn.*
Naturf. 24. 57. 20. *Tab. 2. fig. 24.*
Oliv. Encycl. Méthod. V. 439. 7.
Fonscol. Ann. Sci. Nat. 26. 277.

Chalcis femorata. Fem. *Dalm. Kongl. Vetens. Acad. Handl.*
für är 1820.

Nigra, pilis canis vestita: oculi fusci: ocelli rufo-fusci: antennæ nigræ, apice rufo-fuscae: mesothoracis scutellum apice tubercu-

latum; squamulæ flavæ: abdomen ferè glabrum, sparsè pubescens: pedes rufi; coxæ nigræ; metacoxæ apice rufæ; femora apice supra flava: metafemora subtus dentibus 12 armata; tibiæ basi et apice flavo maculatæ; metatibiæ subtus fuscæ; unguis et pulvilli fusi: alæ hyalinæ; nervi fusi. (Corp. long. 2— $2\frac{1}{2}$ lin.; alar. 4— $4\frac{1}{4}$ lin.)

Var. β. Pro- et mesopedes ferè omnino rufi.

Taken by M. F. de Laporte, near Paris. July; South of France.

Sp. 9. Chal. vicina. Fem. *Nigra, pedibus nigris, femoribus apice tarsisque rufis.*

Nigra, pilis canis vestita: oculi fusi: ocelli rufo-fusi: antennæ nigræ: mesothoracis scutellum inerme; squamulæ flavæ: abdomen ferè glabrum, sparsè pubescens: pedes nigri; trochanteres fusi; pro- et mesofemora apice rufo-flavescens; metafemora apicem versus rufescens, apice supra flavo notata, subtus dentibus 12 armata; pro- et mesotibiæ nigro-fuscæ, apice basi subtusque rufescens; metatibiæ nigræ, supra basim versus et apice flavo maculatae; tarsi pallidè rufi; unguis et pulvilli fusi: alæ subfuscæ; nervi fusi. (Corp. long. $1\frac{3}{4}$ —2 lin.; alar. $3\frac{3}{4}$ —4 lin.)

Taken near Paris, by M. F. de Laporte.

Sp. 10. Chal. parvula. Mas et Fem. *Nigra, pedibus nigris, femoribus apice tarsisque flavis.*

Chalcis minuta. *Dalm. Kongl. Vetens. Acad. Handl. für är 1820.*

Mas. Nigra, pilis canis vestita: oculi fusi: ocelli rufo-fusi: antennæ nigræ, apice rufo-fuscæ: mesothoracis scutellum apice bispinosum; squamulæ flavæ: abdomen ferè glabrum, sparsè pubescens: pedes nigri; trochanteres fusi; femora apice flava; mesofemora subtus dentibus 12 armata; pro- et mesotibiæ nigro-fuscæ, apice basi subtusque flavæ; metatibiæ extus basi et apice flavescentes; tarsi flavi; unguis et pulvilli fusi: alæ hyalinæ; nervi fusi.

Fem. Antennæ paullò breviores et crassiores: scutellum inerme; abdomen longius. (Corp. long. 1— $1\frac{3}{4}$ lin.; alar. 2— $3\frac{3}{4}$ lin.)

Taken near Paris, by M. F. de Laporte. July; South of France.

GENUS IV. HALTICELLA,^f *Spinola.*

Chalcis. . *Fabricius*, *Gmelin*, *Rossi*, *Hubner*, *Olivier*, *Latreille*, *Spinola*.

Halticella. *Spinola*.

Caput mediocre, transversum, thorace vix angustius, anticè ubi insident scapi excavatum : antennæ 13-articulatæ, prope os insertæ ; *maris* subfusciformes, *fem.* subclavatæ, paullò breviores ; articulus 1^{us}. elongatus, 2^{us}. cyathiformis, 3^{us}. minimus, 4^{us}. et 6 sequentes subæquales, magni ; clava triarticulata, conica, articulis 2 præcedentibus brevior : mandibulæ arcuatæ, apice bidentatae ; dentes vix acuti : maxillæ elongatae, apice intùs in lobum quasi articulatum productæ ; palpi 4-articulati, breves ; articulus 1^{us}. medioceris ; 2^{us}. crassior, cyathiformis ; 3^{us}. parvus ; 4^{us}. elongatus, fusiformis, setosus : mentum elongato-ovatum ; palpi 3-articulati, menti apice insertæ, breves ; articulus 2^{us}. minimus : labium rotundum, integrum, anticè ciliatum : thorax ovatus : pectus parvum : prothoracis scutellum mediocre, subquadratum, posticè incisum : mesothoracis sentum magnum ; parapsidum suturæ distinctæ ; scutellum magnum, subrotundum : metathoracis scutellum, paraptera et epimera magna ; illum medio canaliculatum : petiolus brevissimus : *maris* abdomen ovatum, convexum ; segmentum 1^{um}. maximum, abdominis dimidio vix brevius ; 2^{um}. mediocre ; sequentia parva ; subtus abdomen segmenta 6 ventralia conspicua, quorum 1^{um}. et 2^{um}. magna, cætera parva : *fem.* abdomen elongato-ovatum, subtus carinatum ; segmenta ventralia occulta : propedes mediocres ; femora subincrassata ; tibiæ apice spina elongata valida arcuata armatae : mesopedes paullò tenuiores ; spina tibialis multo brevior et gracilior : metapedes coxis femoribusque maximis ; haec subtus dentibus 12 minutis armata ; tibiæ arcuatæ, subtus canaliculatae, apice intùs productæ, acuminatae et spina armatae : alæ *Chaleidis*.

Sp. 1. Hal. pusilla. Mas et Fem. *Nigra*, *femoribus apice flavis*, *tarsis fulvis*, *alis subfuscis*.

Chalcis pusilla. . *Fabr. Mant. Ins.* I. 272. 5. *Ent. Syst.* II. 197. 8. *Syst. Piezat.* 167. 29. *Gmel. Syst. Nat.* I. 5. 2743. 6. *Hübn. Naturf.* 24. 57. 21. *Tab. 2. fig. 25.* *Ross. Faun. Etrusc.* II. 59. 807. *Oliv. Encycl. Méthod.* V. 439. 8.

Halticella pusilla. *Spin. Ann. Mus. Hist. Nat.* Tom. XVII.

^f ἀλτικὸς saltator, κέλλω celeriter moveo. It is usually spelt *Halticella*.

Nigra, punctata, pilis canis vestita : oculi fusi : ocelli rufo-fusi : mesothoracis scutellum inerme ; squamulæ flavæ : abdomen nitidum, glabrum, basi nudum : pedes nigri ; pro- et mesofemora apice fulva ; metafemora apice flava ; pro- et mesotibiæ fuscae, basi, apice et subtus fulvae ; metatibiæ apice et macula basim versus fulvae ; tarsi fulvi ; unguis et pulvilli fusi : alæ subfuscæ, medio obscuriores : nervi fusi. (Corp. long. $1\frac{1}{4}$ — $1\frac{3}{4}$ lin. ; alar. $2\frac{1}{4}$ — $2\frac{3}{4}$ lin.)

Taken near Paris by M. F. de Laporte.

GENUS V. HOCKERIA,^g *De Laporte.*

Chalcis . . Fabricius, Gmelin, Olivier, Latreille.

Halticella. Spinola, Olivier, &c.

Hockeria. De Laporte.

Caput *Halticellæ* : antennæ 13-articulatae, prope os insertæ ; *maris* subfusciformes, corporis dimidio longiores ; *fem.* subclavatae, paullò breviores ; articulus 1^{us}. flagelli dimidii longitudinem ; 2^{us}. cya-thiformis ; 3^{us}. et 8-sequentes subæquales ; 12^{us}. et 13^{us}. minimi, vix conspicui : os *Halticellæ* : thorax ovatus : pectus parvum : prothoracis scutellum magnum, quadratum : mesothoracis scutum magnum ; parapsides benè determinatae ; scutellum magnum, subrotundum ; paraptera et epimera mediocria : metathoracis scutellum maximum, medio canaliculatum : *maris* abdomen sessile, ovatum, convexum ; segmentum 1^{um}. magnum ; sequentia parva, apicem versus longitudine decrescentia ; subtus abdomen segmenta 7 ventralia conspicua, basalia apicalibus longiora : *fem.* abdomen elongato-ovatum, subtus carinatum ; segmentum 1^{um}. maximum, abdominis dimidio paullò brevius ; 2^{um}. mediocre ; 3^{um}. 4^{um}. et 5^{um}. parva ; 6^{um}. latius ; segmenta ventralia vix conspicua : propedes mediocres ; tibiae apice spina armatae ; tarsi breves ; unguis et pulvilli minuti : mesopedes paullò tenuiores : metapedes elongati ; coxae maximæ, trigonæ ; femora magna, ovata, subtus apice dentibus 2 magnis obtusis armata ; tibiæ arcuatæ, subtus canaliculatae, apice latores et spinis 2 armatae : proalæ nervus ordinarius costæ partem brevissimam occupans ; ramulus stigmaticalnis minimus, vix fureatus.

This genus forms the second division of *Chalcis*, in Latreille's *Gen. Crust. et Ins. &c.* Spinola placed it with *Chalcis pusilla* in his genus *Halticella*, which was probably formed on the species belonging to it ; but this being uncertain, I have

^g ὁγκηρὸς tumidus.

adopted De Laporte's generic name. The peculiar form and low insertion of the antennæ, and the very short part of the costa occupied by the ordinary nervures of the superior wings, are characters possessed also by the *Encyrtidæ*.

Sp. 1. Hoc. bispinosa. Fem. *Nigra, pro- et mesotibiis tarsisque omnibus rufis, proalis medio fuscis, albo maculatis.*

Chalcis bispinosa . . Fabr. Syst. Piezat. 166. 28. Fonscol.
Ann. Sci. Nat. 26. 279. 9.

Halticella bispinosa. Spin. Ann. Mus. Hist. Nat. Tom.
XVII. Oliv. Nouv. Dict. d'Hist. Nat.

Nigra, nitida, punctata, vix pilosa: oculi fusci: ocelli rufo-fusci: mesothoracis scutellum apice bispinosum; squamulæ rufo-fuscæ: abdomen nitidissimum, glabrum, acuminatum, subtus rufo-fuscum; pedes nigri; pro- et mesotibiae omnino, et metatibiae apice rufæ: tarsi rufi; unguis et pulvilli fusci: alæ subfuscæ; proalæ medio fuscae, albo bimaculatae; nervi fusci. (Corp. long. 2 lin.; alar. 2 $\frac{3}{4}$ lin.)

Taken near Paris, by M. F. de Laporte.

Sp. 2. Hoc. bifasciata. Fem. *Nigra, tarsis rufis, proalis fusco bifasciatis.*

Chalcis bimaculata. Fonscol. Ann. Sci. Nat. XXVI. 280. 11.

*Nigra, obscura, punctata, pilosa: oculi fusci: ocelli rufo-fusci: antennæ nigræ, graciles, thorace longiores: mesothoracis scutellum inerme; squamulæ nigræ: abdomen *H. bispinosæ* sed brevius, nitidum, glabrum, basi nudum, subtus rufo-fuscum: pedes nigri; femora postica subtus fusca; coxae, tibiæ apice tarsique rufescentes; unguis et pulvilli fusci: alæ subfuscæ; proalarum fasciæ medio connectæ: nervi fusci. (Corp. long. 1—1 $\frac{1}{2}$ lin.; alar. 2—2 $\frac{1}{4}$. lin.)*

Taken near Paris, by M. F. de Laporte.

Sp. 3. Hoc. hetera. Mas. *Nigra, pedibus rufis, metafemoribus et tibiis nigris, mesothoracis scutello integro.*

Nigra, nitida, punctata, pilosa: oculi fusci: ocelli rufo-fusci: antennæ nigræ, gracillimæ, corporis dimidio longiores; scapus perlongus: mesothoracis scutellum inerme; squamulæ rufo-

fuscæ : metathoracis scutellum maximum, apice utrinque produetum : abdomen nitidum, glabrum, subtus apice rufo-fuseum : pedes rufi ; coxae nigrae ; trochanteres fusci ; pro- et mesofemora basi nigra ; metafemora nigra, apice supra rufa ; metatibiæ nigrae, apice rufæ ; tarsi et ungues fusci : alæ subfuscæ ; proalæ medio obscuriores ; nervi fusci. (Corp. long. $1\frac{1}{4}$ lin. ; alar. $2\frac{1}{4}$ lin.)

Taken near Paris, by M. F. de Laporte.

Sp. 4. Hoc. nigra. Mas. *Nigra, metafemoribus tarsisque omnibus rufis.*

Chalcis Dargelasii? *Latr. Hist. Nat. des Crust. &c.* XIII. 221.

Nigra, obscura, punetata, pilosa : oculi fusci : ocelli rufo-fusei : mandibulæ rufæ : antennæ nigrae, thoraci breviores : mesothoracis scutellum inerme ; squamulæ nigro-fuscæ : metathoracis scutellum maximum : abdomen nitidum, glabrum : pedes nigri ; metafemora rufa, basi extus, nonnunquam quoque supra et subtus nigra ; tarsi rufi ; ungues et pulvilli fusci : alæ hyalinæ ; nervi pallidè fusci. (Corp. long. $1\frac{3}{4}$ —2 lin. ; alar. $2\frac{1}{2}$ — $2\frac{3}{4}$ lin.)

Taken near Paris, by M. F. de Laporte.

Sp. 5. Hoc. nigripes. Mas. *Nigra, tarsis rufis, mesothoracis scutello integro.*

Chalcis nigripes. *Fonscol. Ann. Sci. Nat.* XXVI. 280. 10.

Nigra, obseura, punctata, pubescens : oculi fusci : ocelli rufo-fusei : antennæ nigrae : mesothoracis scutellum convexum, integrum ; squamulæ nigrae : metathoracis scutellum maximum, striatum : abdomen nitidum, glabrum, basi nudum : pedes nigri ; metacoxæ nitidissimæ ; trochanteres, ungues et pulvilli fusci : tibiæ apice rufæ : tarsi rufi : alæ subhyalinæ ; nervi fusci, basi pallidiores. (Corp. long. $1\frac{1}{2}$ lin. ; alar. $2\frac{1}{4}$ lin.)

Taken near Paris, by M. F. de Laporte.

Sp. 6. Hoc. rufipes. Mas. *Nigra, pedibus rufis, metafemoribus et tibiis nigris, mesothoracis scutello bispinoso.*

Chalcis rufipes . *Oliv. Encycl. Méthod.* V. 440. 11.

Chalcis clavipes? *Rossi.*

Chalcis armata . *Dalm. Kongl. Vetens. Acad. Handl. für är.* 1820. Var. ?

Cynips armata . *Panz. Faun. Insect. Germ.* 74. 9. Var. ?

Nigra, obscura, punctata, pilosa: oculi fusti: ocelli rufo-fusti: antennæ nigræ, graciles, corporis dimidio longiores: mesothoracis scutellum elevatum, apice spinis duabus longis obtusis armatum; squamulæ nigro-fuscae: abdomen nitidum, glabrum, ferè nudum: pedes rufi; coxæ nigræ; trochanteres fusti; pro- et mesofemora fusco cingulata; metafemora nigra; metatibiæ nigræ, apice rufæ; tarsi et ungues fusti: alæ subfuscæ; proalæ medio obscuriores, macula prope stigma hyalina: nervi fusti. (Corp. long. 1—2 lin.; alar. 2—2 $\frac{3}{4}$ lin.)

Var. β. Pro- et mesotibiæ fusco cingulatae.

Taken near Paris, by M. F. de Laporte; also in England, by Mr. Curtis. *Var. β* is described from an English specimen.

Sp. 7. Hoc. unicolor. Mas. *Nigra, pedibus omnino nigris, alis hyalinis.*

Nigra, obscura, punctata, pilosa: oculi fusti: ocelli rufo-fusti: antennæ nigræ, corporis dimidi longitudinem: mesothoracis scutellum elevatum, apice spinis duabus brevissimis armatum; squamulæ nigræ: abdomen nitidum, glabrum, ferè nudum: pedes nigri: alæ hyalinæ; nervi fusti. (Corp. long. 1 lin.; alar. 1 $\frac{3}{4}$ lin.)

July; South of France.

GENUS VI. NOTASPIS,^h Walker.

Mas.—Caput magnum, transversum, thorace latius, anticè ubi insident scapi excavatum: oculi magni, prominentes, globosi: antennæ 13-articulatae, subclavatae, apice acuminatae, prope os insertae, corporis dimidio breviores; articulus 1^{us}. antennæ triente longior; 2^{us}. mediocris; 3^{us}. minimus; 4^{us}. et 6 sequentes mediocres, subæquales; clava conica, articulis 9^o. et 10^o. longior: thorax ovatus: pectus parvum: prothoracis scutellum mediocre, subquadratum: mesothoracis scutum mediocre; parapsides bene determinatae, magnæ, convexæ; squamulæ maximæ, globosæ; paraptera magna; scutellum maximum, metathoracem abdominisque basim transiens, apice acuminatum: metathorax parvus: abdomen sessile, ovatum, convexum; segmentum 1^{um}. maximum; cætera minima: pro- et mesopedes mediocres, femora subclavata, tarsi graciles; metapedes magni, coxæ trigonæ, femora ovata, tibiæ subtus canaliculatae, tarsi crassi brevesque: alæ *Hockeriæ*.

^h νῶτος dorsum, ἀσπίς clypeus.

Sp. 1. Not. formiciformis. Mas. *Aeneus, antennis fuscis, tarsis fulvis, alis albis.*

Aenea, obscura, punctata, haud pubescens: oculi fusi: ocelli rufo-fusi: antennæ fuscæ; articulus 1^{us}. æneus: thorax punctis magnis profundèque excavatis scaber; latera ferè glabra; squamulæ rufo-fuscæ, nitidæ; mesothoracis scutellum elevatum, subtus apicem unidentatum: abdomen nitidum, glabrum: pedes nigro-ænei; trochanteres fusi; tibiæ apice fulvæ; tarsi fulvi; ungues et pulvilli fusi: alæ albi, iridescentes; nervi pallidi, vix conspicui. (Corp. long. $\frac{4}{5}$ lin.; alar. $1\frac{1}{4}$ lin.)

Taken in St. Vincent's island, by the Rev. Lansdown Guilding.

GENUS VII. DIRHINUS, Dalman.

Chalcis. *Jurine*, Latreille, *Spinola*.

Mas.—Caput magnum, thorace non angustius, multò longius quam latum, anticè inter oculos tuberculis duobus armatum, posticè sub prothorace productum; tubercula lata, obtusa, apice serrata: oculi mediocres, globosi: antennæ 13-articulatae, subclavatae, per longum striatæ; articulus 1^{us}. elongatus; sequentes mediocres; ultimus minimus, vix conspicuus: mandibulæ arcuatæ; una bidentata; altera tridentata: thorax elongato-ovatus: pectus parvum: prothoracis scutellum magnum, subquadratum: mesothoracis scutum angustum; parapsides benè determinatae; scutellum mediocre: metathorax magnus: abdomen ovatum, petiolatum, supra planum, subtus carinatum, apice retusum: segmenta dorsalia subtus abdomen marginem formantia; 1^{um}. s. petiolus crassum; 2^{um}. maximum; cætera minima: segmenta nonnulla ventralia conspicua: tibiæ apice spina armatae: pro- et mesofemora clavata: metapedes magni; coxae elongatae; femora ovata, subtus serrata; tibiæ arcuatæ, subtus canaliculatae; tarsi graciles: proalæ angustæ; nervus ordinarius costam longum occupans; ramulus stigmatical is vix ullus.

This genus is allied to *Cerocephala*, *Spalangia*, &c.

Sp. 1. Dir. cornigerus. Mas. *Ater, genubus tarsisque rufis, alis hyalinis.*

Chalcis cornigera. *Jur. Nouv. Méthod. Hyménopt.* 315. Pl. 13. 47. *Spin. Ins. Lig.* Fascic. 3^{us}. 164. 8.

Nigra, obscura, punctata, pubescens : oculi fusci : ocelli rufo-fusci : antennæ nigræ : caput scabrum : thorax lævior : squamulæ rufo-fuscae : metathorax carinatus, utrinque spinosus : abdomen nitidum, glabrum, basi striatum : pedes nigri ; trochanteres fusci ; pro- et mesofemora apice rufa ; metafemora basi unidentata ; pro- et mesotibiæ basi apiceque rufæ ; tarsi rufi ; ungues et pulvilli fusci : alæ hyalinæ ; proalæ ad costam fuscæ ; nervi fusci. (Corp. long. 2 lin. ; alar. 2 $\frac{1}{4}$ lin.)

Taken near Paris, by M. F. de Laporte.

ART. III.—*Capture of Insects at Burghfield*.—By the Rev.
C. S. BIRD, M.A. F.L.S.

Burghfield Hill House, near Reading, Aug. 1833.

SIR,—Having resided at this place about ten years, and employed my leisure hours in making an Entomological Collection, I think I have ascertained pretty nearly what this locality affords, amongst the more conspicuous insects at least; and, with your permission, I shall be happy to register, in your Magazine, the result of my researches. The country around me is woody, particularly abounding in elms, and my house is close to several copses, containing large, though not old, oaks, &c. ; and at the distance of half a mile I have the range of a heathy common, terminating in fir-groves. There is no chalk, that I am aware of, within six miles.

I am particularly attached to *Lepidoptera*,—probably only because I have been most successful in this order. This success I owe to the use of a lamp to attract moths. During the moonless nights of summer, I sit with a Sinumbra-lamp, and perhaps one or two smaller lamps, placed on a table, close to the window. The moths speedily enter the room, if the weather be warm. I have had a levee of more than a hundred between the hours of ten and twelve. In the spring, too, and autumn, I have been frequently fortunate, though generally having my patience sufficiently tried. In March, for instance, I have taken many specimens of *Biston prodromarius* in one evening ; *Glaea rubricosa*, and *Lytæa leucographa*, have accompanied them. In April and May, *Cucullia fissina*, and *Peridea serrata*, have visited me. When November has arrived,